










# Topic: Vectors

Topic/Skill	Definition/Tips	Example
1. Translation	<p><b>Translate</b> means to <b>move a shape</b>. The shape does not change <b>size</b> or <b>orientation</b>.</p>	
2. Vector Notation	<p>A vector can be written in 3 ways:</p> <p style="text-align: center;"><math>\mathbf{a}</math> or <math>\overrightarrow{AB}</math> or <math>\begin{pmatrix} 1 \\ 3 \end{pmatrix}</math></p>	
3. Column Vector	<p>In a column vector, the <b>top</b> number moves <b>left (-)</b> or <b>right (+)</b> and the <b>bottom</b> number moves <b>up (+)</b> or <b>down (-)</b></p>	<p><math>\begin{pmatrix} 2 \\ 3 \end{pmatrix}</math> means '2 right, 3 up'</p> <p><math>\begin{pmatrix} -1 \\ -5 \end{pmatrix}</math> means '1 left, 5 down'</p>
4. Vector	<p>A <b>vector</b> is a quantity represented by an arrow with both <b>direction</b> and <b>magnitude</b>.</p> <p style="text-align: center;"><math>\overrightarrow{AB} = -\overrightarrow{BA}</math></p>	
5. Magnitude	<p>Magnitude is defined as the <b>length</b> of a vector.</p>	 <div style="border: 1px solid black; padding: 5px; width: fit-content; margin-left: 20px;"> <p>Magnitude (length) can be calculated using Pythagoras Theorem:  <math>3^2 + 4^2 = 25</math>  <math>\sqrt{25} = 5</math></p> </div>
6. Equal Vectors	<p>If two vectors have the <b>same magnitude and direction</b>, they are <b>equal</b>.</p>	

<p>7. Parallel Vectors</p>	<p><b>Parallel</b> vectors are <b>multiples</b> of each other.</p>	<p><math>2\mathbf{a}+\mathbf{b}</math> and <math>4\mathbf{a}+2\mathbf{b}</math> are parallel as they are multiple of each other.</p> 
<p>8. Collinear Vectors</p>	<p><b>Collinear</b> vectors are vectors that are on the <b>same line</b>. To show that two vectors are <b>collinear</b>, show that one vector is a <b>multiple</b> of the other (parallel) <b>AND</b> that both vectors <b>share a point</b>.</p>	
<p>9. Resultant Vector</p>	<p>The <b>resultant</b> vector is the vector that results from <b>adding</b> two or more vectors together.</p> <p>The resultant can also be shown by <b>lining up the head</b> of one vector with the <b>tail</b> of the other.</p>	<p>if <math>\underline{\mathbf{a}} = \begin{pmatrix} 4 \\ 4 \end{pmatrix}</math> and <math>\underline{\mathbf{b}} = \begin{pmatrix} 2 \\ -2 \end{pmatrix}</math></p> <p>then <math>\underline{\mathbf{a}} + \underline{\mathbf{b}} = \begin{pmatrix} 4 \\ 4 \end{pmatrix} + \begin{pmatrix} 2 \\ -2 \end{pmatrix} = \begin{pmatrix} 6 \\ 2 \end{pmatrix}</math></p> 
<p>10. Scalar of a Vector</p>	<p>A <b>scalar</b> is the <b>number</b> we multiply a vector by.</p>	 <p>Example:</p> $3\mathbf{a} + 2\mathbf{b} =$ $= 3\begin{pmatrix} 2 \\ 1 \end{pmatrix} + 2\begin{pmatrix} 4 \\ -1 \end{pmatrix}$ $= \begin{pmatrix} 6 \\ 3 \end{pmatrix} + \begin{pmatrix} 8 \\ -2 \end{pmatrix}$ $= \begin{pmatrix} 14 \\ 1 \end{pmatrix}$

## 11. Vector Geometry




$$\vec{OA} = a \quad \vec{AO} = -a$$

$$\vec{OB} = b \quad \vec{BO} = -b$$

$$\vec{AB} = \vec{AO} + \vec{OB} = -a + b = b - a$$

$$\vec{BA} = \vec{BO} + \vec{OA} = -b + a = a - b$$

**Example 1:**  $X$  is the midpoint of  $AB$ . Find  $\vec{OX}$   
**Answer:** Draw  $X$  on the original diagram



Now build up a journey.

You could use  $\vec{OX} = \vec{OA} + \frac{1}{2}\vec{AB}$ .

This will give:  $\vec{OX} = a + \frac{1}{2}(b - a)$ .

This will simplify to  $\frac{1}{2}a + \frac{1}{2}b$  or  $\frac{1}{2}(a + b)$